



PCT/EP200 4 / 0 0 4 7 9 9



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 21 JUN 2004

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

**CERTIFIED COPY OF
PRIORITY DOCUMENT**

Signed

Dated 20 April 2004

BEST AVAILABLE COPY



The
Patent
Office


10JUN03 E813554-4 002898
P01/7700 0.00-0313249.5

The Patent Office

Cardiff Road
Newport
Gwent NP10 8QQ

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



1.	Your reference	C4313 (C)/TC		
2.	Patent application number (The Patent Office will fill in this part)	0313249.5		- 9 JUN 2003
3.	Full name, address and postcode of the or of each applicant (underline all surnames)	UNILEVER PLC UNILEVER HOUSE, BLACKFRIARS LONDON, EC4P 4BQ		
	Patents ADP number (if you know it)	50426956002		
	If the applicant is a corporate body, give the country/state of its incorporation	UNITED KINGDOM		
4.	Title of the invention	BLEACHING COMPOSITION		
5.	Name of your agent (if you have one)	ELLIOTT Peter William		
	"Address for Service" in the United Kingdom to which all correspondence should be sent (including the postcode)	PATENT DEPARTMENT, UNILEVER PLC COLWORTH HOUSE, SHARNBROOK BEDFORD, MK44 1LQ		
	Patents ADP number (if you know it)	6573927001		
6.	If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)
7.	If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application		Date of filing (day/month/year)
8.	Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. See note (d))	YES		

Patents Form 1/77

9. ☒ Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	19
Claim(s)	3
Abstract	1
Drawing(s)	

g

10. If you are also filing any of the following, state how many against each item.

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

1 /

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature(s) *Leonie Watkinson* Date: 09/06/03

Leonie WATKINSON, Authorised Signatory

12. Name and daytime telephone number of person to contact in the United Kingdom

Trudi Clark (01234) 222360

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

- 1. -

BLEACHING COMPOSITION

FIELD OF INVENTION

This invention relates to the enhancement of bleaching
5 compositions that are substantially devoid of peroxy species.

BACKGROUND OF INVENTION

The use of bleaching catalysts for stain removal has been
10 developed over recent years. The recent discovery that
some catalysts are capable of bleaching effectively in
the absence of an added peroxy source has recently
become the focus of some interest, for example:
WO9965905; WO0012667; WO0012808; WO0029537, and,
15 WO0060045.

The shelf life of a product may be regarded as the period
of time over which the product may be stored whilst
retaining its required quality. A satisfactory shelf
20 life is in many instances a crucial factor for the
success of a commercial product. A product with a short
shelf life generally dictates that the product is made in
small batches and is rapidly sold to the consumer. It is
also a concern to the owners of a brand with a short
25 shelf life that the consumer uses the product within the
shelf life otherwise the consumer may be inclined to
change to a similar product of another brand. In
contrast a similar product with a long shelf life may be
made in larger batches, held as stock for a longer period
30 of time and the period of time that a consumer stores the

- 2 -

product is not of a great concern to the owners of a particular brand.

It is an object of the present invention to provide an
5 air bleaching composition that has improved storage properties.

SUMMARY OF INVENTION

We have found that some components degrade per se and/or
10 reduce the activity of bleaching catalysts over a period of time. We have found that by carefully selecting certain components the stability of a bleaching composition, which is substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach
15 system, is improved. The present invention has particular utility in commercial liquid bleaching compositions.

The present invention provides a bleaching composition comprising:

20

(a) an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous
25 bleaching medium substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system,

(b) between 0.001 to 3 wt/wt % of a perfume composition
30 said perfume composition comprising at least 0.01 wt % of an ketonic perfume, and

(c) an antioxidant in the range from 0.0001 to 20 wt/wt %, .

(d) the balance carriers and adjunct ingredients to 100 wt/wt % of the total bleaching composition.

The term "substantially devoid of a peroxygen bleach or a peroxy-based or peroxy-generating bleach system" should be construed within spirit of the invention. It is preferred that the composition has as low a content of peroxy species present as possible. It is preferred that the bleaching formulation contains less than 1 % wt/wt total concentration of peracid or hydrogen peroxide or source thereof, preferably the bleaching formulation contains less than 0.3 % wt/wt total concentration of peracid or hydrogen peroxide or source thereof, most preferably the bleaching composition is devoid of peracid or hydrogen peroxide or source thereof. In addition, it is preferred that the presence of alkyl hydroperoxides is kept to a minimum in a bleaching composition comprising the ligand or complex of the present invention.

The present invention extends to a method of bleaching a substrate/textile with a composition of the present invention. The method comprising the steps of treating a substrate with the bleaching composition in an aqueous environment, rinsing the substrate and drying the substrate.

The present invention also extends to a commercial package together with instructions for its use.


DETAILED DESCRIPTION OF THE INVENTION

ANTIOXIDANT

The bleaching compositions of the present invention will
5 comprise an effective amount of the antioxidant. An
effective amount of an antioxidant is in the range 0.001
to 20 wt/wt % depending upon the nature of the
antioxidant and subsidiary purpose of the antioxidant,
for example as a carrier or solvent. Preferably the
10 antioxidant is present in the range from 0.001 to 2 wt/wt
%. When a phenolic antioxidant is present it is
preferred that the phenolic antioxidant present in the
range from 0.0001 to 3 % wt % of the composition. When an
amine antioxidant it is present it is preferred that the
15 phenolic antioxidant present in the range from 0.0001 to
20 % wt % of the composition.

Anti-oxidants are substances as described in Kirk-Othmers
(Vol 3, pg 424) and in Uhlmans Encyclopedia (Vol.3, pg
20 91).


One class of anti-oxidants suitable for use in the
present invention is alkylated phenols having the general
formula:



25 wherein R is C1-C22 linear or branched alkyl, preferably
methyl or branched C3-C6 alkyl; C3-C6 alkoxy, preferably

methoxy; R1 is a C3-C6 branched alkyl, preferably tert-butyl; x is 1 or 2. Hindered phenolic compounds are preferred as antioxidant.

- 5 Another class of anti-oxidants suitable for use in the present invention is a benzofuran or benzopyran derivative having the formula:



- wherein R1 and R2 are each independently alkyl or R1 and R2 can be taken together to form a C5-C6 cyclic hydrocarbonyl moiety; B is absent or CH2; R4 is C1-C6 alkyl; R5 is hydrogen or -C(O)R3 wherein R3 is hydrogen or C1-C19 alkyl; R6 is C1-C6 alkyl; R7 is hydrogen or C1-C6 alkyl; X is -CH2OH, or -CH2A wherein A is a nitrogen comprising unit, phenyl, or substituted phenyl. Preferred nitrogen comprising A units include amino, pyrrolidino, piperidino, morpholino, piperazino, and mixtures thereof.

- Other suitable antioxidants are found as follows. A derivative of α -tocopherol, beta-tocopherol, gamma-tocopherol, delta-tocopherol, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox™).

- Anti-oxidants/radical scavengers such as ascorbic acid (vitamin C) and its salts, tocopherol (vitamin E), tocopherol sorbate, other esters of tocopherol, butylated hydroxy benzoic acids and their salts, gallic acid and its alkyl esters, especially propyl gallate, uric acid

and its salts and alkyl esters, sorbic acid and its salts, the ascorbyl esters of fatty acids, amines (e.g., N,N-diethylhydroxylamine, amino-guanidine, amine alcohols), sulfhydryl compounds (e.g., glutathione), and
5 dihydroxy fumaric acid and its salts may be used.

Non-limiting examples of anti-oxidants suitable for use in the present invention include phenols *inter alia* 2,6-di-tert-butylphenol, 2,6-di-tert-butyl-4-methylphenol,
10 mixtures of 2 and 3- tert-butyl-4-methoxyphenol, and other ingredients including include propyl gallate, tert-butylhydroquinone, benzoic acid derivatives such as methoxy benzoic acid, methylbenzoic acid, dichloro benzoic acid, dimethyl benzoic acid, 5-hydroxy-2,2,4,6,7-
15 pentamethyl-2,3-dihydro-1-benzofuran-3-one, 5-hydroxy-3-methylene-2,2,4,6,7-pentamethyl-2,3-dihydro-benzofuran, 5-benzyloxy-3-hydroxymethyl-2,2,4,6,7-pentamethyl-2,3-dihydro-1-benzofuran, 3-hydroxymethyl-5-methoxy-
2,2,4,6,7-pentamethyl-2,3-dihydro-1-benzofuran, vitamin
20 C(ascorbic acid), and Ethoxyquine (1,2-dihydro-6-ethoxy-2,2,4-trimethylchinolin) marketed under the name Raluquin™ by the company Raschig™.


Preferred antioxidants for use herein include 2,6-di-
25 tert-butyl hydroxy toluene (BHT), α -tocopherol, hydroquinone, 2,2,4-trimethyl-1,2-dihydroquinoline, 2,6-di-tert-butyl hydroquinone, 2-tert-butyl hydroquinone, tert-butyl-hydroxy anisole, lignosulphonic acid and salt thereof, benzoic acid and derivatives thereof, like
30 alkoxyated benzoic acids, as for example, trimethoxy benzoic acid (TMBA), toluic acid, catechol, t-butyl

catechol, benzylamine, amine alcohols, 1,1,3-tris(2-methyl-4-hydroxy-5-t-butylphenyl) butane, N-propyl-gallate or mixtures thereof and highly preferred is di-tert-butyl hydroxy toluene. Of the amine alcohols 2-amino-2-methyl-1-propanol, tri-ethanol amine, tri-methanol amine, mono-ethanol amine, diethanol amine, are preferred.


Mixtures of antioxidants may be use and in particular mixtures that have synergic antioxidant effects as found in, for example; WO02/072746.

KETONIC PURFUMES


The bleaching composition of the present invention comprises between 0.001 to 3 wt/wt % of a perfume composition, preferably between 0.1 to 2 wt/wt % of a perfume composition. The perfume composition may be 100% ketonic perfume but generally the perfume composition is a complex mixture of perfumes of other differing perfume classifications, for example terpenes and aldehydes; in this regard, the perfume composition comprises at least 0.01 wt % of a ketonic perfume. At higher levels of ketonic perfume the greater the importance of the antioxidant, for example at least 0.1, 1.0 wt % and 5 wt % of a ketonic perfume component of the perfume composition. The present invention has particular utility with, but not limited to, the following ketonic perfume components.




pulegone



vertofix coeur




veloutone



Alpha-methylionone

5



damascenone

THE BLEACH CATALYST

The bleach catalyst per se may be selected from a wide
 10 range of transition metal complexes of organic molecules
 (ligands). In typical washing compositions the level of
 the organic substance is such that the in-use level is
 from 0.05 μM to 50 mM, with preferred in-use levels for
 domestic laundry operations falling in the range 1 to 100
 15 μM . Higher levels may be desired and applied in
 industrial textile bleaching processes.

Suitable organic molecules (ligands) for forming complexes and complexes thereof are found, for example in:


GB 9906474.3; GB 9907714.1; GB 98309168.7, GB

5 98309169.5; GB 9027415.0 and GB 9907713.3; DE 19755493; EP 999050; WO-A-9534628; EP-A-458379; EP 0909809; United States Patent 4,728,455; WO-A-98/39098; WO-A-98/39406, WO 9748787, WO 0029537; WO 0052124, and WO0060045 the complexes and organic molecule (ligand) precursors of
10 which are herein incorporated by reference. An example of a preferred catalyst is a transition metal complex of MeN₄Py ligand (N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane).

15 The ligand forms a complex with one or more transition metals, in the latter case for example as a dinuclear complex. Suitable transition metals include for example: manganese in oxidation states II-V, iron II-V, copper I-III, cobalt I-III, titanium II-IV, tungsten IV-VI,
20 vanadium II-V and molybdenum II-VI.

An example of a preferred catalyst is a monomer ligand or transition metal catalyst thereof of a ligand having the formula (I):

25




wherein each R is independently selected from: hydrogen, F, Cl, Br, hydroxyl, C1-C4-alkylo-, -NH-CO-H, -NH-CO-C1-C4-alkyl, -NH₂, -NH-C1-C4-alkyl, and C1-C4-alkyl;

R1 and R2 are independently selected from:

- 5 C1-C4-alkyl,
- C6-C10-aryl, and,
- a group containing a heteroatom capable of coordinating to a transition metal, wherein at least one of R1 and R2 is the group containing the heteroatom;
- 10 R3 and R4 are independently selected from hydrogen, C1-C8 alkyl, C1-C8-alkyl-O-C1-C8-alkyl, C1-C8-alkyl-O-C6-C10-aryl, C6-C10-aryl, C1-C8-hydroxyalkyl, and -(CH₂)_nC(O)OR₅ wherein R₅ is independently selected from: hydrogen, C1-C4-alkyl, n is from 0 to 4, and mixtures thereof; and,
- 15 X is selected from C=O, -[C(R₆)₂]_Y- wherein Y is from 0 to 3 each R₆ is independently selected from hydrogen, hydroxyl, C1-C4-alkoxy and C1-C4-alkyl.

The transition metal complex preferably is of the general
20 formula (AI):



in which:

- 25 M represents a metal selected from Mn(II) - (III) - (IV) - (V), Cu(I) - (II) - (III), Fe (II) - (III) - (IV) - (V), Co(I) - (II) - (III), Ti(II) - (III) - (IV), V(II) - (III) - (IV) - (V), Mo(II) - (III) - (IV) - (V) - (VI) and W(IV) - (V) - (VI), preferably from Fe(II) - (III) - (IV) - (V);

L represents the ligand, preferably N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, or its protonated or deprotonated analogue;

X represents a coordinating species selected from
5 any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner;

Y represents any non-coordinated counter ion;

a represents an integer from 1 to 10;

10 k represents an integer from 1 to 10;

n represents zero or an integer from 1 to 10;

m represents zero or an integer from 1 to 20.

BALANCE CARRIERS AND ADJUNCT INGREDIENTS

15

These are generally surfactants, builders, foam agents, anti-foam agents, solvents, and enzymes. The use and amounts of these components are such that the bleaching composition performs depending upon economics.

20 environmental factors and use of the bleaching composition.

The air bleach catalyst may be used in a detergent composition specifically suited for stain bleaching

25 purposes, and this constitutes a second aspect of the invention. To that extent, the composition comprises a surfactant and optionally other conventional detergent ingredients. The invention in its second aspect provides an enzymatic detergent composition which comprises from
30 0.1 - 50 % by weight, based on the total detergent composition, of one or more surfactants. This surfactant

system may in turn comprise 0 - 95 % by weight of one or more anionic surfactants and 5 to 100 % by weight of one or more nonionic surfactants. The surfactant system may additionally contain amphoteric or zwitterionic detergent compounds, but this is not normally desired owing to their relatively high cost. The enzymatic detergent composition according to the invention will generally be used as a dilution in water of about 0.05 to 2%.

- 10 In general, the nonionic and anionic surfactants of the surfactant system may be chosen from the surfactants described "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch, Interscience 1958, in the current edition of
- 15 "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners Company or in "Tenside-Taschenbuch", H. Stache, 2nd Edn., Carl Hauser Verlag, 1981.
- 20 Suitable nonionic detergent compounds which may be used include, in particular, the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example, aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially
- 25 ethylene oxide either alone or with propylene oxide. Specific nonionic detergent compounds are C₆-C₂₂ alkyl phenol-ethylene oxide condensates, generally 5 to 25 EO, i.e. 5 to 25 units of ethylene oxide per molecule, and the condensation products of aliphatic C₈-C₁₈ primary or
- 30 secondary linear or branched alcohols with ethylene oxide, generally 5 to 40 EO.

Suitable anionic detergent compounds which may be used are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl radicals containing from about 8 to about 22 carbon atoms, the term alkyl being used to include the alkyl portion of higher acyl radicals. Examples of suitable synthetic anionic detergent compounds are sodium and potassium alkyl sulphates, especially those obtained by sulphating higher C₈-C₁₈ alcohols, produced for example from tallow or coconut oil, sodium and potassium alkyl C₉-C₂₀ benzene sulphonates, particularly sodium linear secondary alkyl C₁₀-C₁₅ benzene sulphonates; and sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil and synthetic alcohols derived from petroleum. The preferred anionic detergent compounds are sodium C₁₁-C₁₅ alkyl benzene sulphonates and sodium C₁₂-C₁₈ alkyl sulphates. Also applicable are surfactants such as those described in EP-A-328 177 (Unilever), which show resistance to salting-out, the alkyl polyglycoside surfactants described in EP-A-070 074, and alkyl monoglycosides.

Preferred surfactant systems are mixtures of anionic with nonionic detergent active materials, in particular the groups and examples of anionic and nonionic surfactants pointed out in EP-A-346 995 (Unilever). Especially preferred is surfactant system that is a mixture of an alkali metal salt of a C₁₆-C₁₈ primary alcohol sulphate together with a C₁₂-C₁₅ primary alcohol 3-7 EO ethoxylate.

The nonionic detergent is preferably present in amounts greater than 10%, e.g. 25-90% by weight of the surfactant system. Anionic surfactants can be present for example in amounts in the range from about 5% to about 40% by weight of the surfactant system.

One skilled in the art will appreciate that some adventitious peroxy species may be in the composition nevertheless it is most preferred that the bleaching composition of the present invention has less than 1%, preferably less than 0.1%, most preferably less than 0.01%, of a peroxy species present. These adventitious peroxy are predominantly alkyl hydroperoxides formed by autoxidation of the surfactants.

The detergent composition may take any suitable physical form, such as a powder, granular composition, tumble dryer sheet, tablets, a paste or an anhydrous gel.

The composition may contain additional enzymes as found in WO 01/00768 A1 page 15, line 25 to page 19, line 29, the contents of which are herein incorporated by reference.

Builders, polymers and other enzymes as optional ingredients may also be present as found in WO0060045.

Suitable detergency builders as optional ingredients may also be present as found in WO0034427.

The composition of the present invention may be used for laundry cleaning, hard surface cleaning (including cleaning of lavatories, kitchen work surfaces, floors, mechanical ware washing etc.). As is generally known in the art, bleaching compositions are also employed in waste-water treatment, pulp bleaching during the manufacture of paper, leather manufacture, dye transfer inhibition, food processing, starch bleaching, sterilisation, whitening in oral hygiene preparations and/or contact lens disinfection.

In the context of the present invention, bleaching should be understood as relating generally to the decolourisation of stains or of other materials attached to or associated with a substrate. However, it is envisaged that the present invention can be applied where a requirement is the removal and/or neutralisation by an oxidative bleaching reaction of malodours or other undesirable components attached to or otherwise associated with a substrate. Furthermore, in the context of the present invention bleaching is to be understood as being restricted to any bleaching mechanism or process that does not require the presence of light or activation by light.

LIQUID FORMULATION

The present invention has particular utility for liquid formulations because in contrast to a solid heterogeneous mixture in a liquid formulation the contact between individual components are more intimate and hence more

susceptible to degradation due to interaction of components.

There are many commercial liquid formulations for
5 detergents and rinse conditioners or other liquid
products that may be enhanced by conferring a bleaching
ability to the liquid formulation. As will be evident to
one skilled in the art the present invention is
applicable to known liquid formulations and liquid
10 formulations to be developed.

The level of the catalyst in a commercial bleaching
composition is from 0.0001 to 0.6 wt/wt %, preferably
0.001 to 0.15 wt/wt %, most preferably 0.01 to 0.1 wt/wt
15 %. We have found that the level of catalyst is optimum
between 0.03 to 0.09 wt/wt % in the commercial bleaching
composition.

The present invention extends to both isotropic and
20 complex liquid compositions and formulations a brief
discussion of which follows. Some isotropic formulations
are termed 'micro-emulsion' liquids that are clear and
thermodynamically stable over a specified temperature
range. The 'micro-emulsion' formulation may be water in
25 oil, or oil in water emulsions. Some liquid formulations
are macro-emulsions that are not clear and isotropic.
Emulsions are considered meta-stable. Concentrated, clear
compositions containing fabric softening actives have
been disclosed in WO 98/08924 and WO 98/4799, both
30 Procter & Gamble. Such compositions comprise bio-
degradable fabric conditioners. However, both disclose

compositions comprising water miscible solvents that do not form water-in-oil micro-emulsions. Clear fabric conditioning compositions have also been disclosed in EP 730023 (Colgate Palmolive), WO 96/19552 (Colgate Palmolive), WO 96/33800 (Witco Co.), WO 97/03170 (Procter & Gamble), WO 97/03172 (Procter & Gamble), WO 97/03169 (Procter & Gamble), US 5492636 (Quest Int.) and US 5427697 (Procter & Gamble). Liquid formulations of the present invention may contain for example; monoethoxy quats; AQAs and bis-AQAs; cationic amides; cationic esters; amino/diamino quats; glucamide; amine oxides; ethoxylated polyethyleneimines; enhancement polymers of the form linear amine based polymers, e.g. bis-hexamethylenetriamine; polyamines e.g. TETA, TEPA or PEI polymers.

The liquid may be contained within a sachet as found in WO02/068577.

The following is an example of a liquid bleaching composition to which an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen may be added together further antioxidant if required. The perfume composition as found in the following example comprising at least 0.01 wt % of an ketonic perfume.

Ingredient	Wt%
Nonionic surfactant	26.6
Monopropylene glycol	5.5
Pigment premix	0.017
Glycerol	21.36
Monoethanolamine	7.56
Oleic fatty acid	13.10
Water	Up to 100
Linear alkyl benzene sulfonate	20.1
Perfume	1.6
Protease Enzyme	1.0

The following are further examples of commercial liquid formulations that the present invention may be incorporated into: Wisk™ liquid USA, 1999, OMO™ liquid NL, 1999, OMO-liquido™ Brazil, 1999, and Rinse conditioner (Robijn™ - NL). In this regard, that catalyst is added and the level of antioxidant adjusted together with the perfume composition.

The following is a further example of a commercial liquid formulation that the present invention may be incorporated into by adding the catalyst and the appropriate level of antioxidant and perfume composition. The commercial liquid formulation has a pH of 7.

Sodium citrate:	3.2 %
Polypropylene glycol:	4.75 %
LAS-acid:	5.6 %
NI 25 9 EO:	6.6 %
LES (anionic surfactant):	10.5 %
Borax:	2.30 %
Sorbitol:	3.35 %
Alcosperce 725:	0.30 %
Coconut fatty acid:	0.73 %
monoethanolamine:	0.20 %
fluorescer:	0.125 %
savinase/lipex	
perfume/dye	

It is most preferred that when the catalyst is in a liquid composition, the liquid composition has a pH of 7 or below irrespective of whether perfumes and/or antioxidants are present or not.

We claim:

1. A bleaching composition comprising:

- 5 (a) an organic substance which forms a complex with a transition metal for bleaching a substrate with atmospheric oxygen, the bleaching composition upon addition to an aqueous medium providing an aqueous bleaching medium substantially devoid of a peroxygen
10 bleach or a peroxy-based or peroxy-generating bleach system,
- (b) between 0.001 to 3 wt/wt % of a perfume composition said perfume composition comprising at least 0.01 wt % of
15 an ketonic perfume, and
- (c) an antioxidant in the range from 0.0001 to 20 wt/wt %,
- 20 (d) the balance carriers and adjunct ingredients to 100 wt/wt % of the total bleaching composition.

2. A bleaching composition according claim 1, wherein the antioxidant is selected from:

25

(i) an phenolic antioxidant, the phenolic antioxidant present in the range 0.0001 to 3 % wt %; and,

(ii) an amine antioxidant, the amine antioxidant in the
30 range from 0.0001 to 20%.

3. A bleaching composition according claim 2, wherein the amine antioxidant is an amine alcohol.

4. A bleaching composition according to claim 3,
5 wherein the amine alcohol is selected from the group consisting of: 2-amino-2-methyl-1-propanol, tri-ethanol amine, tri-methanol amine, mono-ethanol amine, diethanol amine, and methylantranilate.

10 5. A bleaching composition according to claim 2, wherein the antioxidant is a hindered phenol.

6. A bleaching composition according to claim 5,
wherein the antioxidant is selected from the group
15 consisting of: 2, 6-di-tert-butyl hydroxy toluene, α -tocopherol, Ethoxyquine and 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid, and lignosulphonic acid.

7. A bleaching composition according to claim 6,
20 wherein the antioxidant is 2, 6-di-tert-butyl hydroxy toluene.

8. A bleaching composition according to claim 1,
wherein the antioxidant is ascorbic acid.

25

9. A bleaching composition according to claim 5 to 8,
wherein the antioxidant is present in the bleaching
composition in the range from 0.001 to 2 wt %.

30 10. A bleaching composition according to any preceding claim, wherein the bleaching composition is a liquid.

11. A bleaching composition according to any preceding claim, comprising between 0.1 to 2 wt/wt % of a perfume composition.

5 12. A bleaching composition according to any preceding claim, wherein said perfume composition comprises at least 0.1 wt % of a ketonic perfume.

10 13. A bleaching composition according to claim 12, wherein said perfume composition comprises at least 1.0 wt % of a ketonic perfume.

15 14. A bleaching composition according to claim 13, wherein the perfume composition comprises at least 5 wt % of a ketonic perfume.

20 15. A bleaching composition according to any one of claims 1 to 9 and 11 to 14, wherein the bleaching composition is a solid bleaching composition.

25 16. A bleaching composition according to any preceding claim, wherein the ketonic perfume is selected from the group consisting of: pulegone, vertofix coeur, veloutone, alpha-methylionone, and damascenone.

30 17. A method of bleaching a textile stain, comprising the steps of treating a substrate with the bleaching composition as defined in any preceding claim in an aqueous environment, rinsing the substrate and drying the substrate.

Abstract of the Invention

The present invention concerns the preservation of a perfume components is a bleaching composition. The bleaching composition is substantially devoid of a .
5 peroxygen bleach or a peroxy-based or peroxy-generating bleach system.

THIS PAGE BLANK (USPTO)

BEST AVAILABLE COPY

EP 04/004799

